

Algebraic Scaling of the Kondo Temperature in a Luttinger Liquid

Philip Phillips

*Loomis Laboratory of Physics
University of Illinois at Urbana-Champaign
1100 W.Green St., Urbana, IL, 61801-3080*

I show here how the algebraic scaling of the Kondo temperature in a Luttinger liquid can be understood simply from the condition for the existence of the singlet (or triplet) bound state.

PACS numbers:72.10.Fk,71.45.-d, 72.15.Nj

In a recent letter Furusaki and Nagaosa (FN) [1] (and Lee and Toner (LT) [2] previously) addressed the problem of the Kondo effect in a Luttinger liquid. One of the key results of FN is that the Kondo temperature scales algebraically as $T_k \propto J^{2\eta}$, with J the Kondo exchange coupling, $\eta = \frac{\pi v_F}{g_2}$, v_F the Fermi velocity and g_2 the coupling constant for forward electron scattering. This result is valid strictly in the limit of weak Coulomb interactions, $\frac{g_2}{2\pi v_F} \ll 1$. Earlier LT bosonized the extended Hubbard model and obtained as an exponent $\eta = \frac{1}{2-K_\rho-K_\sigma^{-1}}$, the spin-density-wave exponent [2]. K_σ and K_ρ are the spin and charge coupling constants for the Luttinger liquid. Although this exponent is valid in the weak-coupling regime, it does not reduce to the simple form predicted by FN. FN pointed out that because bosonization does not preserve the $SU(2)$ symmetry inherent in the Kondo model, exponents obtained from this procedure will be in error [1]. However, the consensus does seem to be that the Kondo effect in a Luttinger liquid is fundamentally distinct from the standard Kondo problem in a Fermi liquid in which the bound singlet state forms only when $J < 0$ and T_k decays exponentially with the Kondo coupling as $T_k \propto e^{\left(\frac{1}{n(\epsilon_F)J}\right)}$.

What I want to show here is that the algebraic scaling of the Kondo temperature in a Luttinger liquid obtained by FN has a simple origin and can be predicted without recourse to renormalization group analyses. Further, the expression I derive is valid regardless of the strength of the electron interactions. My starting point is the Fermi liquid problem in which the zeroth-order Kondo singlet ground state can be written [3] as

$$|\Psi\rangle = \sum_{k>k_F} \alpha_k (a_{k\uparrow}^\dagger \chi_\downarrow - a_{k\downarrow}^\dagger \chi_\uparrow) |F\rangle \quad (1)$$

where $|F\rangle$ is the full Fermi sea $|F\rangle = \prod_{k<k_F} a_{k\uparrow}^\dagger a_{k\downarrow}^\dagger |0\rangle$, α_k is an expansion coefficient and χ_s is the spin eigenstate of the local magnetic impurity. The expansion coefficient is easily shown to be given by $\alpha_k = \langle F | \chi_\downarrow^\dagger a_{k\uparrow} - a_{k\downarrow} \chi_\uparrow^\dagger | \Psi \rangle =$

$\langle F | b_k | \Psi \rangle$. The time evolution of α_k can be obtained from the Heisenberg equations of motion $i\dot{\alpha}_k = \langle F | [b_k, H] | \Psi \rangle$, where H is the standard Kondo Hamiltonian. If we write $\alpha_k(t) = e^{-iEt} \alpha_k(t=0)$, we find that the evolution equation reduces to the stability condition

$$1 = -J_o \sum_k' \frac{1}{E - \epsilon_k} \quad (2)$$

with $J_o = \frac{J}{N}$. The sum is strictly above the Fermi surface and the ϵ_k are the energies of the free conduction states. Eq. (2) is the standard Cooper-like equation, the solution of which yields the energy (below the continuum) at which the bound singlet Kondo state forms. Conversion of the sum in (2) to an integral introduces the single particle density of states. For the Fermi liquid problem, this quantity can be treated as a constant and the right-hand side of (2) is easily integrated to yield a logarithm. The exponential dependence of the bound state energy on the Kondo coupling J_o follows immediately.

In the corresponding Luttinger problem, a similar derivation can be followed because the bosonization procedure generates a set of non-interacting psuedo-particles [4]. In this case, however, the single particle density of states is not a constant. Rather, it scales as a power law and vanishes at the Fermi level. Below the Kondo temperature, strong coupling between the impurity and the electron spin located at the origin effectively breaks the correlated 1-d electron system into two semi-infinite Luttinger liquids [1]. The Kondo impurity appears as a boundary condition at one end of the chain. For the $SU(2)$ symmetric case in which $K_\sigma = 1$, the appropriate density of states for a semi-infinite Luttinger liquid is $n(\epsilon) \propto |\epsilon|^{\frac{1}{g}-1}$ in the vicinity of the Fermi level [5]. The coupling constant g is the dimensional conductance in the Luttinger liquid, $g = \frac{1}{\sqrt{1 + \frac{g_2}{\pi v_F}}}$. For repulsive interactions, $g_2 > 0$ and consequently, $g < 1$. If we now substitute this form for the density of states [6] in Eq. (2) and convert the sum to an integral, we find immediately that the integral has an algebraic singularity that scales as $E_b^{-(\frac{1}{g}-1)}$. The stability condition for the singlet state now reduces to

$$1 = \text{const.} J_o E_b^{-(\frac{1}{g}-1)} \quad (3)$$

which implies that

$$E_b \propto J_o^{\frac{1}{\frac{1}{g}-1}} = k_B T_k. \quad (4)$$

Let us now analyze this expression in the weak-coupling regime. In this regime g can be expanded in a power series in $\frac{g_2}{\pi v_F} \ll 1$. We find that the leading power law dependence, $T_k \propto J_o^{\left(\frac{2\pi v_F}{g_2}\right)}$, is precisely the result of FN.

In the strong-coupling regime, $T_k \propto J_o^{\sqrt{\frac{\pi v_F}{g_2}}}$ [7]. The procedure of FN is not easily amenable to a strong-coupling analysis. We find then that this simple procedure based on the analysis of the stability condition for the Kondo bound state predicts algebraic scaling of the Kondo temperature that recovers the weak coupling result. It is the power law dependence of the pseudo-particle density of states for a semi-infinite Luttinger liquid that leads to this behaviour. The physical picture that this derivation suggests is that the pseudo-particles in the Luttinger liquid form the screening cloud at the magnetic impurity in direct analogy to the quasi-particles in the Fermi liquid problem. Of course this picture changes dramatically if back-scattering is included [8]. As FN pointed out, in the presence of backscattering, the two semi-infinite pieces of the Luttinger liquid become connected and the scaling behaviour presented here breaks down. Away from half-filling, however, back scattering as well as umklapp processes are not relevant along the Luttinger fixed line. Hence, away from half-filling, the analysis presented here should hold true.

ACKNOWLEDGMENTS

This work is supported in part by the NSF and the ACS Petroleum Research Fund. I thank Yi Wan, Nancy Sandler, and A.-H. Castro Neto for insightful remarks.

-
- [1] A. Furusaki and N. Nagaosa, Phys. Rev. Lett. **72**, 892 (1994).
 - [2] D. H. Lee and J. Toner, Phys. Rev. Lett. **69**, 3378 (1992).
 - [3] K. Yosida, Phys. Rev. **147**, 223 (1966).
 - [4] F. D. M. Haldane, J. Phys. C **14**, 2585 (1981).
 - [5] C. L. Kane and M. P. A. Fisher, Phys. Rev. Lett. **68**, 1220 (1992).
 - [6] This derivation is in the spirit of the interlayer tunneling argument used by S. Chakravarty and P. W. Anderson, Phys. Rev. Lett. **72** 3859 (1994).
 - [7] The results presented here can be trivially extended to the triplet ground state, which also exhibits a Kondo effect in a Luttinger liquid. In Eq. 1, a plus sign should be used and $J \rightarrow -J$.
 - [8] A. Luther and V. J. Emery, Phys. Rev. Lett. **33**, 589 (1974).